

74 Rec'd PCT/FTO 14 AUG 1997
08/894246

11 Rec'd PCT/PTO 14 AUG 1997

FORM PTO-1390
(REV. 5-93)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

EX95001-US

U.S. APPLICATION NO.
(If known. See 37 C.F.R. 1.5)

INTERNATIONAL APPLICATION NO.

PCT/FR96/00218

INTERNATIONAL FILING DATE

12 February 1996

PRIORITY DATE CLAIMED

14 February 1995

TITLE OF INVENTION

MEDICINAL COMBINATION USEFUL FOR IN VIVO EXOGENIC TRANSFECTION AND EXPRESSION

APPLICANT(S) FOR DO/EO/US

Michel PERRICAUDET, Martin LEE, Lucienne CHATENOUD, Hedi HADDADA, Jean-François BACH, and
Michelle WEBB

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)), and Power of Attorney, unsigned.
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:

11. ☐ An information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

CERTIFICATION UNDER 37 CFR 1.10

GB840711938 US

"Express Mail" Mailing Number

14 August 1997

Date of Deposit

I hereby certify that this paper (along with any referred to as being attached or enclosed) is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box PCT, Assistant Commissioner for Patents, Washington, D.C. 20231, Attn. EO/US

Paula L. Dickey
(Type or print name of person mailing paper)Paula L. Dickey
(Signature of person mailing paper)

U.S. APPLICATION NO. (If Known, see C.F.R. 1.5)	INTERNATIONAL APPLICATION NO. PCT/FR96/00218	ATTORNEY'S DOCKET NUMBER EX95001-US
--	---	--


17. <input checked="" type="checkbox"/> The following fees are submitted: Basic National Fee (37 CFR 1.492(a)(1)-(5): Search Report has been prepared by the EPO or JPO.....\$ 910.00 International preliminary examination fee paid to USPTO (37 CFR 1.482)\$ 700.00 No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2))..\$ 770.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid USPTO.....\$1040.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4).....\$ 96.00 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>	CALCULATIONS PTO use only																				
Surcharge of \$130.00 for furnishing the oath or declaration later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(e)).	\$ 910.00																				
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 20%;">Claims</th> <th style="width: 20%;">Number Filed</th> <th style="width: 20%;">Number Extra</th> <th style="width: 20%;">Rate</th> <th style="width: 20%;"></th> </tr> <tr> <td>Total Claims</td> <td>31 -20 =</td> <td>11</td> <td>X \$ 22.00</td> <td>\$ 242.00</td> </tr> <tr> <td>Independent Claims</td> <td>2 - 3 =</td> <td>0</td> <td>X \$ 80.00</td> <td>\$ 0.00</td> </tr> <tr> <td colspan="4">Multiple dependent claim(s) (if applicable)</td> <td>+ \$260.00</td> </tr> </table>	Claims	Number Filed	Number Extra	Rate		Total Claims	31 -20 =	11	X \$ 22.00	\$ 242.00	Independent Claims	2 - 3 =	0	X \$ 80.00	\$ 0.00	Multiple dependent claim(s) (if applicable)				+ \$260.00	
Claims	Number Filed	Number Extra	Rate																		
Total Claims	31 -20 =	11	X \$ 22.00	\$ 242.00																	
Independent Claims	2 - 3 =	0	X \$ 80.00	\$ 0.00																	
Multiple dependent claim(s) (if applicable)				+ \$260.00																	
TOTAL OF ABOVE CALCULATIONS =	\$ 1152.00																				
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28).	\$																				
SUBTOTAL =	\$ 1152.00																				
Processing fee of \$130.00 for furnishing the English translation later the [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).	\$																				
TOTAL NATIONAL FEE =	\$ 1152.00																				
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +	\$																				
TOTAL FEES ENCLOSED =	\$ 1152.00																				
	Amount to be refunded \$																				
	charged \$																				

a. ☐ A check in the amount of \$_____ to cover the above fee is enclosed.

b. ☒ Please charge my Deposit Account No. 18-1982 in the amount of \$1152.00 to cover the above fees.
 A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 18-1982. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO: Paul F. Fehlner, Esquire Rhone-Poulenc Rorer Inc. Legal-Patents, #3C43 P.O. Box 5093 Collegeville, PA 19426-0997 Telephone: (610) 454-3839 Facsimile: (610) 454-3808	<div style="text-align: center;">  Signature </div> <div style="text-align: center;"> Paul F. Fehlner, Ph.D. Name </div> <div style="text-align: center;"> 35,135 Registration Number </div> <div style="text-align: center;"> Date <u>14 August 1997</u> </div>
---	---

74 Rec'd PCT/PTO 14 AUG 1997

08/894246

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of: Perricaudet et al. Group Art Unit:
Serial No.: To Be Assigned Examiner:
U.S. National Stage of PCT/FR96/000218
Filed: Concurrently Herewith
For: Medicinal Combination Useful For In Vivo Exogenic Transfection and
Expression
To: The Honorable Commissioner of Patents and Trademarks
Washington, D.C. 20231

CERTIFICATE OF MAILING (37 CFR § 1.10)

GB840711938 US
"Express Mail" Mailing Number

14 August 1997
Date of Deposit

I hereby certify that this paper (along with any referred to as being attached or enclosed) is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box PCT, Assistant Commissioner for Patents, Washington, D.C. 20231, Attn. EO/US.

Paula L. Dickey
(Type or print name of person mailing paper)

Paula L. Dickey
(Signature of person mailing paper)

PRELIMINARY AMENDMENT

Please enter the following amendment to the English Translation of the International Application before examining this application.

In the Claims

Please cancel claims 1-25 without prejudice.

Please add the following new claims:

-- 26. A composition comprising an immunosuppressive agent and a recombinant adenovirus whose genome comprises a first recombinant DNA containing a therapeutic gene and a second recombinant DNA containing an immunoprotective gene.

27. The composition according to claim 26, wherein the immunosuppressive agent is selected from the group consisting of cyclosporin,

FK506, azathioprine, corticosteroids and monoclonal or polyclonal antibodies that are able to inactivate immune molecules or induce destruction of the immune cells carrying these molecules.

28. The composition according to claim 27, wherein the antibody is selected from among the anti-CD4, -CD2, -CD3, -CD8, -CD28, -B7, -ICAM-1 and -LFA-1 antibodies and CTLA4Ig.

29. The composition according to claim 26, wherein the therapeutic gene encodes a therapeutic protein.

30. The composition according to claim 26, wherein the therapeutic gene encodes a therapeutic RNA.

31. The composition according to claim 26, wherein the immunoprotective gene is a gene whose product acts on the activity of the major histocompatibility complex (MHC) or on the activity of the cytokines.

32. The composition according to claim 31, wherein the immunoprotective gene is a gene whose product at least partially inhibits expression of the MHC proteins or antigen presentation.

33. The composition according to claim 26, wherein the immunoprotective gene is selected from the group consisting of a gene for gp19k of adenovirus, the ICP47 gene of herpes virus, and the UL18 gene of cytomegalovirus.

34. The composition according to claim 26, wherein the two recombinant DNAs of the adenovirus genome constitute a single transcriptional entity.

35. The composition according to claim 26, wherein the two recombinant DNAs each include an identical transcriptional promoter.

36. The composition according to claim 35, wherein the two recombinant DNAs are inserted in the same orientation.

37. The composition according to claim 26, wherein the two recombinant DNAs are inserted into the same region of the adenovirus genome.

38. The composition according to claim 37, wherein the two recombinant DNAs are inserted within the E1, E3 or E4 regions.

39. The composition according to claim 26, wherein the two recombinant DNAs are inserted into different sites in the adenovirus genome.

40. The composition according to claim 39, wherein one of the recombinant DNAs is inserted within the E1 region and the other within the E3 or E4 region.

41. The composition according to claim 26, wherein the adenovirus is a defective recombinant adenovirus which encompasses the ITR sequences and a sequence permitting encapsidation and which carries a deletion of all or part of the E1 and E4 genes.

42. The composition according to claim 26, wherein the adenovirus concerned is an adenovirus from whose genome all or part of the E1, E3, L5 and E4 genes have been deleted.

43. A method for expression of a therapeutic gene from an adenovirus comprising consecutively or simultaneously administering an immunosuppressive agent and a recombinant adenovirus whose genome comprises a first recombinant DNA containing a therapeutic gene and a second recombinant DNA containing an immunoprotective gene, to a subject.

44. The method according to claim 43, wherein the recombinant adenovirus is administered in vivo.

45. The method according to claim 43, wherein the immunosuppressive agent is selected from the group consisting of cyclosporin, FK506, azathioprine, corticosteroids and monoclonal or polyclonal antibodies that are able to inactivate immune molecules or induce destruction of the immune cells carrying these molecules.

46. The method according to claim 45, wherein the antibody is selected from among the anti-CD4, -CD2, -CD3, -CD8, -CD28, -B7, -ICAM-1 and -LFA-1 antibodies and CTLA4Ig.

47. The method according to claim 43, wherein the therapeutic gene encodes a therapeutic protein.

48. The method according to claim 43, wherein the therapeutic gene encodes a therapeutic RNA.

49. The method according to claim 43, wherein the immunoprotective gene is a gene whose product acts on the activity of the major histocompatibility complex (MHC) or on the activity of the cytokines.

50. The method according to claim 49, wherein the immunoprotective gene is a gene whose product at least partially inhibits expression of the MHC proteins or antigen presentation.

51. The method according to claim 43, wherein the immunoprotective gene is selected from the group consisting of a gene for gp19k of adenovirus, the ICP47 gene of herpes virus, and the UL18 gene of cytomegalovirus.

52. The method according to claim 43, wherein the two recombinant DNAs of the adenovirus genome constitute a single transcriptional entity.

53. The method according to claim 43, wherein the two recombinant DNAs each include an identical transcriptional promoter.

54. The method according to claim 53, wherein the two recombinant DNAs are inserted in the same orientation.

55. The method according to claim 43, wherein the immuno-suppressive agent is injected both before and after injection of the adenovirus.

56. The method according to claim 43, wherein the immuno-suppressive agent and the recombinant adenovirus are injected simultaneously. --


REMARKS

Claims 1-25 have been cancelled and rewritten as new claims 26-56 in order to conform with US patent practice. In addition, new method claims 43-56 are supported at page 1, lines 3-9, page 5, lines 7-11, and page 8, lines 1-2. No new matter has been added.

Rhône-Poulenc Rorer Inc.
Legal-Patents #3C43
P.O. Box 5093
Collegeville, PA 19426-0997
Telephone: (610) 454-3839
Facsimile: (610) 454-3808

Dated: 8/14/97

Respectfully submitted,



Paul F. Fehlner, Ph.D.
Attorney for Applicants
Registration No. 35,135

WO 96/25177

1

PCT/FR96/00218

MEDICINAL COMBINATION USEFUL FOR IN VIVO EXOGENIC TRANSFECTION
AND EXPRESSION

The present invention relates to the field of gene therapy and in particular to the use of adenovirus for expressing a therapeutic gene of interest. It relates, more specifically, to a novel method for treating pathologies of genetic origin, which method is based on the combined use of two types of therapeutic agents.

Gene therapy consists in correcting a deficiency or an anomaly (mutation, aberrant expression, etc.) by introducing genetic information into the affected cell or organ. This genetic information can be introduced either in vitro or ex vivo into a cell which has been removed from the organ, with the modified cell then being reintroduced into the organism, or else directly in vivo into the appropriate tissue. In this second case, a variety of different physical techniques exist for transfection, including the use of viruses as vectors. In this respect, a variety of different viruses have been tested for their ability to infect particular cell populations. These viruses include, in particular, retroviruses (RSV, HMS, MMS, etc.), the HSV virus, adeno-associated viruses and adenoviruses.

Among these viruses, the adenoviruses exhibit some properties which are favourable in relation to use in gene therapy. They have a rather broad host spectrum,

are capable of infecting quiescent cells and do not integrate into the genome of the infected cell. The adenoviruses are viruses which contain linear double-stranded DNA of about 36 kb in size. Their genome encompasses, in particular, an inverted repeat sequence (ITR) at their end, an encapsidation sequence, early genes and late genes (cf. Figure 1). The principal early genes are the genes E1 (E1a and E1b), E2, E3 and E4. The principal late genes are the genes L1 to L5.

In view of the adenovirus properties mentioned above, these viruses have already been used for transferring genes in vivo. To this end, different adenovirus-derived vectors have been prepared which incorporate a variety of different genes (β -gal, OTC, α -IAT, cytokines, etc.). In each of these constructs, the adenovirus was modified in such a way as to render it incapable of replicating in the infected cell. Thus, the constructs which are described in the prior art are adenoviruses from which the E1 (E1a and/or E1b) and, possibly, E3 regions have been deleted, with a heterologous DNA sequence being inserted in their stead (Levrero et al., Gene 101 (1991) 195; Gosh-Choudhury et al., Gene 50 (1986) 161).

However, as in the case for all known viruses, administration of a wild-type virus (Routes et al., J. Virol. 65 (1991) 1450) or of a recombinant virus which is defective for replication (Yang et al., PNAS (1994) 4407) induces a substantial immune response.

The primary aim of the immune system is the integrity of the individual or the integrity of "self". It leads to the elimination of infectious agents and the rejection of transplants and tumours without, however, these powerful defence mechanisms of the organism turning against it and giving rise to autoimmune diseases. This state of non-response with regard to "self" antigens when foreign antigens are eliminated is defined as a state of physiological tolerance. In order to eliminate foreign agents, the immune system develops two types of mechanisms. The first is the production of specific antibodies by the B lymphocytes; this is termed humoral immunity. These antibodies fix the antigen and either inactivate it or eliminate it from the organism. The second defence mechanism involves cellular immunity and employs T lymphocytes, among these the cytotoxic T lymphocytes which carry a specific receptor for the antigen in question. Recognition of the antigen by the T receptor necessitates the latter being expressed in association with proteins which are encoded via the genes of the major histocompatibility complex or class I and class II MHC.

Consequently, this immune response, which is developed against the infected cells, constitutes a major obstacle to the use of viral vectors in gene therapy since (i) by inducing destruction of the infected cells it limits the period during which the therapeutic gene is expressed and hence the therapeutic effect, (ii) it

induces, in parallel, a substantial inflammatory response, and (iii) it brings about rapid elimination of the infected cells after repeated injections. It will be understood that the amplitude of this immune response against infected cells varies according to the nature of the organ which sustains the injection and according to the method of injection which is employed. Thus, expression of the β -galactosidase encoded by a recombinant adenovirus which is administered into the muscle of immunocompetent mice is reduced to minimum levels 40 days after the injection (Kass-Eisler et al., PNAS 90 (1993) 11498). In the same way, the expression of genes which have been transfected into the liver using adenoviruses is significantly reduced in the 10 days following the injection (Yang Y et al. 1994 immunity 1 433-442) and expression of factor IX which was transferred using adenovirus into the hepatocytes of haemophilic dogs disappeared at 100 days after the injection (Kay et al. PNAS 91 (1994) 2353).

From the point of view of exploiting vectors derived from adenoviruses for the purpose of gene therapy, it therefore seems necessary to control the immune response which is developed against them or against the cells which they are infecting.

From the above, it follows that activation of the immune system first of all requires recognition by the system of elements which are foreign to the organism (non-self or modified self) such as vectors derived from

adenoviruses, which would normally be destroyed. In recent years, immunointervention strategies have been developed whose aim is to create a "permissive" immune environment, that is to say induce a state of tolerance with regard to predefined foreign antigens.

It is precisely at this level that the present invention intervenes. The invention is directed towards preventing the rapid elimination of the adenoviruses from the infected cells and hence towards prolonging, in a consistent manner, the in vivo expression of the therapeutic gene which they are carrying.

Recently, the Applicant has demonstrated that the co-expression of certain genes in the infected cells is able to induce an immunoprotective effect and thus enable the vectors and/or the infected cells to evade the immune system. The Applicant has, in particular, developed adenoviruses in which expression of a gene of therapeutic importance is coupled to that of an immunoprotective gene (FR No. 94 12346). This gene can, in particular, be a gene whose product acts on the activity of the major histocompatibility complex (MHC) or on the activity of the cytokines, thereby making it possible to reduce considerably, if not suppress, any immune reaction against the vector or the infected cells. These gene products at least partially inhibit expression of the MHC proteins or presentation of the antigen, advantageously resulting in a significant reduction of the immune reaction against the vector or the infected

cells, and hence a prolonged therapeutic effect.

Unexpectedly, the Applicant demonstrated that it was possible significantly to prolong, over time, the therapeutic effect of such a vector by associating it with an immunosuppressant. Elimination of the vector in question and/or destruction of the infected cells, by the immune system, is/are found to be retarded over time by a period which is markedly greater than that which might have been expected by the simple juxtaposition of the immunoprotective effects of the said vector and the immunosuppressant. Advantageously, the medicinal combination, which is a subject of the present invention, induces a phenomenon of "pseudo-inertia" of the immune system, which phenomenon favours expression in the long term of a therapeutic gene.

Within the meaning of the invention, immunosuppressant indicates any compound which is able to inhibit, wholly or in part, at least one immune signalling pathway. In general, immunosuppressants are routinely used in transplantation, with the aim of preventing allograft rejection, and in the treatment of certain autoimmune diseases. The products which are customarily used are either chemical immunosuppressants such as corticosteroids, azathioprine, cyclosporin or FK506, or biological immunosuppressants such as polyclonal or monoclonal antibodies. The first category of immunosuppressants, and among these cyclosporin and FK506, in particular, exert a substantial inhibitory

effect on the production of cytokines, such as interleukin 2, which play an essential role in the differentiation and proliferation of the lymphocytic cells. Unfortunately, for this type of immunosuppressant
5 to be effective, they have to be administered continuously, something which sooner or later runs into the problem of their toxicity. Thus, azzathioprine is potentially myelotoxic while cyclosporin is nephrotoxic and can also bring about hypertension or neurological
10 disorders.

As regards the antibodies, more particularly, these are antibodies which are directed against the lymphoid cells of the immune system. The first antibody which was used as an immunosuppressant is anti-CD3, which
15 is directed against the T lymphocytes. Its target is the one of the polypeptide chains of the CD3 molecule which forms the receptor for the T cell antigen. There then follows a functional inactivation of the CD3+ T cells which are recognized by the antibody. As regards the
20 problem which is of interest in the present case, administration of an immunosuppressant of this type together with that of a recombinant adenovirus containing a therapeutic gene would be in a position to block the immune reaction of the host with regard to the viral
25 vector and/or its products which are expressed on the surface of the infected cells. Anti-CD4, -CD2, -CD8, -CD28, -B7, -ICAM-1 and -LFA-1 antibodies can be used on the same principle.

The Applicants have has now developed a novel method of treatment which is particularly efficient in substantially delaying, if not inhibiting, the reaction of the immune system without raising any toxicity
5 problem.

More specifically, the present invention ensues from the demonstration of a particularly substantial synergistic effect which is associated with the combined use of a recombinant adenovirus, in which expression of
10 a gene of therapeutic importance is coupled to that of an immunoprotective gene, such as previously described, and of at least one immunosuppressive agent.

The present invention therefore relates, initially, to a medicinal combination of at least one
15 immunosuppressive agent and at least one recombinant adenovirus whose genome comprises a first recombinant DNA containing a therapeutic gene and a second recombinant DNA containing an immunoprotective gene, for consecutive, intermittent and/or simultaneous use over time, which can
20 be used for exogenous transfections in vivo and/or ex vivo.

As indicated above, the invention is based, in particular, on the demonstration of a synergistic effect between the activity of the immunosuppressive agent and
25 the effect of the expressed immunoprotective gene on the expression of the therapeutic gene.

This combined use makes it possible to achieve a therapeutic effect which is markedly prolonged and

advantageously requires doses which are significantly reduced, in particular as regards their content of immunosuppressive agent.

As indicated further below, the two components of the combined treatment of the present invention can be used consecutively, intermittently and/or simultaneously over time. Preferably, the immunosuppressive agent is injected before and after injection of the adenovirus. According to this method of implementing the present invention, the administration of the immunosuppressant can be spaced out over time and, more preferably, be repeated regularly. In this particular case, the two components are packaged separately. When administration takes place simultaneously, they can be mixed as required before being administered together or, on the other hand, they can be administered simultaneously but separately. In particular, the routes by which the two agents are administered can be different.

According to the present invention, any compound which is able to inhibit, wholly or in part, at least one immune signalling pathway can be used as the immunosuppressive agent. The compound can be selected, in particular, from cyclosporin, FK506, azathioprine, corticosteroids and any monoclonal or polyclonal antibody. Use is preferably made of antibodies which are able to inactivate immune molecules or induce destruction of the immune cells carrying these molecules. Anti-CD4, -CD3, -CD2, -CD8, -CD28, -B7, -ICAM-1 and -LFA-1

antibodies can, in particular, be used as antibodies. Use can also be made of hybrid molecules such as CTLA4Ig, a protein fusion between the CTLA-4 molecule (a homologue of CD28) and an immunoglobulin. The GlFc site of this molecule is found to be able to inhibit activation of the T cells by binding to the B7 molecule (D. J; Lenschow; Science, 257, 789, 1992). It is obvious that the scope of the present invention is in no way limited to the immunosuppressants enumerated above. These immunosuppressants can be employed in isolation or in combination.

The recombinant DNAs which are present in the genome of the adenovirus which is employed in accordance with the present invention are DNA fragments which contain the gene (therapeutic or immunoprotective) under consideration and, where appropriate, signals which enable it to be expressed, and which are constructed in vitro and then inserted into the genome of the adenovirus. The recombinant DNAs which are used within the scope of the present invention can be complementary DNAs (cDNAs), genomic DNAs (gDNAs), or hybrid constructs which consist, for example, of a cDNA in which one or more introns is/are inserted. They can also be synthetic or semisynthetic sequences. These DNAs can be of human, animal, vegetable, bacterial, viral, etc. origin. Use is particularly advantageously made of cDNAs or of gDNAs.

Any gene which encodes a product having a therapeutic effect may be mentioned as a therapeutic gene which can be used for constructing the vectors of the

present invention. The product which is thus encoded can be a protein, a peptide, an RNA, etc.

A protein product can be homologous with regard to the target cell (that is, it can be a product which is normally expressed within the target cell when the latter is not exhibiting any pathology). In this case, expression of a protein makes it possible, for example, to compensate for insufficient expression in the cell or for expression of a protein which is inactive or weakly active due to a modification, or even to overexpress said protein. The therapeutic gene can also encode a mutant of a cell protein, which mutant has an increased stability, a modified activity, etc. The protein product can also be heterologous with regard to the target cell. In this case, an expressed protein can, for example, supplement or provide an activity which is deficient in the cell, thereby permitting the latter to resist a pathology, or else stimulate an immune response.

Those therapeutic protein products within the meaning of the present invention which may more specifically be mentioned are enzymes, blood derivatives, hormones, interleukins, interferons, TNF, etc. (FR 9203120), growth factors, neurotransmitters or their precursors or enzymes for synthesizing them, trophic factors: BDNF, CNTF, NGF, IGF, GMF, aFGF, bFGF, NT3, NT5, HARP/pleiotrophin, etc.; apolipoproteins: ApoAI, ApoAIV, ApoE, etc. (FR 93 05125), dystrophin or a minidystrophin (FR 9111947), the CFTR protein associated with

mucoviscidosis, tumour-suppressing genes: p53, Rb, RplA, DCC, k-rev, etc. (FR 93 04745), genes encoding for factors involved in coagulation: factors VII, VIII and IX, genes intervening in DNA repair, etc.

5 As indicated above, the therapeutic gene can also be an antisense gene or sequence whose expression in the target cell makes it possible to control the expression of genes or the transcription of cellular mRNAs. Such sequences can, for example, be transcribed in
10 the target cell into RNAs which are complementary to cellular mRNAs and thereby block translation of the latter into protein, in accordance with the technique described in Patent EP 140 308. Antisense sequences also include sequences encoding ribozymes, which are able
15 selectively to destroy target RNAs (EP 321 201).

 The therapeutic genes can be of human, animal, vegetable, bacterial, viral, etc. origin. They can be obtained by any technique known to the person skilled in the art and, in particular, by screening libraries, by
20 chemical synthesis or else by mixed methods including chemical or enzymic modification of sequences obtained by screening libraries.

 The immunoprotective gene which is used within the scope of the present invention can be of different
25 types. As previously explained, it is a gene whose product acts on the activity of the major histocompatibility complex (MHC) or on the activity of the cytokines. It is preferably a gene whose product at

least partially inhibits expression of the MHC proteins or antigen presentation. As preferred examples, mention may be made of certain genes contained in the adenovirus E3 region, the herpes virus gene ICP47 or the
5 cytomegalovirus gene UL18.

The E3 region of the adenovirus genome contains different reading frames which, by means of alternative splicing, give rise to different proteins. Among these, the Gp19k (or E3-19k) protein is a glycosylated
10 transmembrane protein which is located in the membrane of the endoplasmic reticulum (RE). This protein encompasses a luminal domain which binds MHC-1 molecules and a C-terminal cytoplasmic end which is able to bind microtubules (or tubulin), which has the effect of
15 anchoring the gp19k protein in the RE membrane. Gp19k is thus able to prevent expression of the MHC-1 molecules at the surface of the cells by interacting with the molecules and sequestering them within the RE. However, protein gp19k is weakly expressed by adenoviruses in the
20 absence of viral replication. Furthermore, expression of gp19k is also dependent on a splicing taking place. Introduction of a recombinant DNA which contains a (preferably cDNA) sequence encoding gp19k into the vectors of the invention enables the expression of said protein to
25 be controlled and optimized. In particular, the use of constitutive promoters and suppression of the other reading frames enables expression of this protein to be strongly increased and freedom to be achieved from

dependence on viral replication and the presence of inducing elements. This makes it possible, particularly advantageously, to considerably diminish lysis of the infected cells by the CTL and thus to increase and
5 prolong the in vivo production of the therapeutic gene.

Other proteins encoded by the E3 region of the adenovirus genome, such as the 10.4k and 14.5k proteins, exhibit certain properties which are attractive with regard to incorporating these genes into the vectors of
10 the invention.

The ICP47 gene of herpes simplex virus represents another immunoprotective gene which is particularly attractive within the meaning of the present invention. Cells which are infected by herpes simplex
15 virus exhibit resistance to lysis induced by CTLs. It has been demonstrated that the ICP47 gene, which can reduce expression of MHC-I molecules at the surface of cells, was able to confer this resistance. Incorporation of the ICP47 gene into a recombinant DNA according to the
20 invention also enables the recombinant viruses of the invention to evade the immune system.

The UL18 gene of cytomegalovirus represents another preferred example of an immunoprotective gene according to the invention. The product of the UL18 gene
25 is able to bind β 2-microglobulin (Brown et al. Nature 347 (1990) 770). β 2-Microglobulin is one of the chains of MHC-I molecules. Incorporation of the UL18 gene into a recombinant DNA according to the invention thus makes it

possible to decrease the number of functional $\beta 2$ -microglobulin molecules in cells infected by the viruses of the invention and therefore to decrease the ability of these cells to produce MHC-I molecules which are complete and functional. This type of construct therefore enables the infected cells to be protected from lysis by CTLs.

As indicated above, the immunoprotective gene which is used within the scope of the present invention is, in another preferred embodiment, a gene whose product inhibits the activity or the signalling pathways of cytokines. The cytokines represent a family of secreted proteins which act as signal molecules for the immune system. They can attract cells of the immune system, activate them and induce them to proliferate, and can even act directly on the infected cells in order to kill them.

Among the genes whose product affects the activity or the signalling pathways of the cytokines, mention may be made of the genes which are involved in the synthesis of the cytokines or whose product is able to sequester cytokines, antagonize their activity or interfere with the intercellular signalling pathways. Preferred examples which may be cited are, in particular, the BCRF1 gene of Epstein Barr virus, the crmA and crmB genes of cowpox virus, the B15R and B18R genes of vaccinia virus, the US28 gene of cytomegalovirus, and the E3-14.7, E3-10.4 and E3-14.5 genes of adenovirus.

The B15R gene of vaccinia virus encodes a

soluble protein which is able to bind interleukin- 1β (the secreted form of interleukin-1) and thereby prevent this cytokine from binding to its cellular receptors. Thus, interleukin-1 is one of the first cytokines to be produced in response to an antigenic attack and it plays a very important role in the signalling of the immune system at the beginning of the infection. The feasibility of incorporating the B15R gene into a vector according to the invention advantageously makes it possible to reduce the activity of IL- 1β , in particular on the activation of the immune cells, and, therefore, provide local protection of the cells which are infected with the viruses of the invention from a significant immune response. Genes which are homologous to the B15R gene, such as the gene of cowpox virus, can also be employed.

In the same way, the B18R gene of vaccinia virus encodes a protein which is homologous to the receptor for interleukin-6. This gene, or any functional homologue, can also be used in the vectors of the invention in order to inhibit binding of interleukin-6 to its cell receptor and thus to reduce the immune response locally.

The crmB gene of cowpox virus can also advantageously be used in a similar fashion. Thus, this gene encodes a secreted protein which is able to bind TNF and to compete with the TNF receptors at cell surfaces. This gene therefore makes it possible, in the viruses of the invention, to locally decrease the concentration of

active TNF which is able to destroy the infected cells. Other genes which encode proteins which are able to bind TNF and at least partially inhibit its binding to its receptors can also be employed.

5 The crmA gene of cowpox virus encodes a protein which has a protease-inhibiting activity of the serpin type and which is able to inhibit the synthesis of interleukin-1 β . This gene can therefore be used in order locally to decrease the concentration of interleukin-1
10 and thus to reduce development of the immune and inflammatory responses.

 The BCRF1 gene of Epstein Barr virus encodes an analogue of interleukin 10. The product of this gene is a cytokine which is able to decrease the immune response
15 and to alter its specificity while inducing proliferation of B lymphocytes.

 The US28 gene of cytomegalovirus encodes a protein which is homologous to the receptor for macrophage inflammatory protein 1 α (MIP-1 α). This protein
20 is therefore able to compete with the receptors for MIP and therefore to inhibit its activity locally.

 The product of the E3-147, E3-10.4 and E3-14.5 genes of adenovirus is able to block transmission of the intercellular signal which is mediated by certain
25 cytokines. When the cytokines bind to their receptor at the surface of an infected cell, a signal is transmitted to the nucleus in order to induce cell death or stop protein synthesis. This is particularly the case for

tumour necrosis factor (TNF). Incorporation of the E3-14.7, E3-10.4 and/or E3-14.5 genes into a recombinant DNA according to the invention for the purpose of expressing them constitutively or in a regulated manner enables
5 intercell signalling which is induced by TNF to be blocked and thus cells which are infected with the recombinant viruses according to the invention to be protected from the toxic effects of this cytokine.

A local and transitory inhibition can be
10 particularly advantageous. This can be obtained, in particular, by the choice of specific expression signals (cytokine-dependent promoters, for example) as indicated below.

It will be understood that other genes which
15 are homologous or which have similar functional properties can be used to construct the vectors of the invention. These different genes can be obtained by any technique which is known to the person skilled in the art and, in particular, by screening libraries, by chemical
20 synthesis or else by mixed methods including chemical or enzymic modification of sequences obtained by screening libraries. Furthermore, these different genes can be employed alone or in combination(s).

Insertion of the genes under consideration in
25 the form of recombinant DNAs according to the invention provides greater flexibility in the construction of the adenoviruses and enables expression of said genes to be controlled more effectively.

Thus, the recombinant DNAs (and therefore the two genes of interest) which are incorporated into the adenoviral vectors according to the present invention can be organized in different ways.

5 They can, first of all, be inserted into the same site in the adenovirus genome or into different, selected sites. In particular, the recombinant DNAs can be inserted, at least in part, into the E1, E3 and/or E4 regions of the adenovirus genome to replace or supplement
10 viral sequences.

 Preferably, the recombinant DNAs are inserted, at least in part, within the E1, E3 or E4 regions of the adenovirus genome. When they are inserted into two different sites, preference is given, within the scope of
15 the invention, to using the E1 and E3 regions or E1 and E4 regions. Thus, as the examples demonstrate, this organization enables the two genes to be expressed at an elevated level without interfering with each other. Advantageously, the recombinant DNAs are inserted in
20 place of viral sequences.

 These recombinant DNAs can then each include a transcriptional promoter which is identical or different. This configuration enables higher levels of expression to be achieved and provides improved control of the
25 expression of the genes. In this case, the two genes can be inserted in the same orientation or in opposite orientations.

 They can also constitute a single

transcriptional entity. In this configuration, the two recombinant DNAs are contiguous and positioned such that the two genes are under the control of a single promoter and give rise to a single premessenger RNA. This arrangement is advantageous since it enables a single transcriptional promoter to be used.

Finally, the use of recombinant DNAs according to the invention makes it possible to employ transcriptional promoters of different types and, in particular, promoters which are strong or weak, regulated or constitutive, tissue-specific or ubiquitous, etc.

The choice of the expression signals and the respective positions of the DNA recombinants is particularly important as regards obtaining an elevated expression of the therapeutic gene and a significant immunoprotective effect.

A particularly preferred embodiment of the present invention employs a defective adenovirus which includes a first recombinant DNA, containing a therapeutic gene, and a second recombinant DNA, containing an immunoprotective gene, in which virus the two recombinant DNAs are inserted within the E1 region.

A particularly preferred embodiment of the present invention employs a defective adenovirus which includes a first recombinant DNA, which contains a therapeutic gene and which is inserted within the E1 region, and a second recombinant DNA, which contains an immunoprotective gene and which is inserted within the E3

region.

As indicated above, the adenoviruses of the present invention are defective, that is they are unable to replicate autonomously in the target cell. Generally, the genome of the defective adenoviruses according to the present invention therefore lacks at least the sequences which are required for replicating said virus in the infected cell. These regions can be eliminated (in whole or in part), rendered non-functional, or substituted by other sequences and, in particular, by the therapeutic genes. The defective character of the adenoviruses of the invention is an important feature since it ensures that the vectors of the invention are not disseminated following administration.

In a preferred embodiment, the adenoviruses of the invention encompass ITR sequences and an encapsidation sequence, and possess a deletion of all or part of the E1 gene.

The inverted repeat (ITR) sequences represent the origin of replication of the adenoviruses. They are located at the 3' and 5' ends of the viral genome (cf. Figure 1), from where they can easily be isolated using standard molecular biological techniques known to the person skilled in the art. The nucleotide sequence of the ITR sequences of the human adenoviruses (in particular serotypes Ad2 and Ad5) is described in the literature, as are those of the canine adenoviruses (in particular CAV1 and CAV2). In the case of the Ad5 adenovirus, for

example, the left-hand ITR sequence corresponds to the region encompassing nucleotides 1 to 103 of the genome.

The encapsidation sequence (also termed Psi sequence) is required for encapsidating the viral DNA.

5 This region must, therefore, be present to enable defective recombinant adenoviruses according to the invention to be prepared. In the adenovirus genome, the encapsidation sequence is located between the left-hand (5') ITR and the E1 gene (cf. Figure 1). It can either be
10 isolated or synthesized artificially using standard molecular biological techniques. The nucleotide sequence of the encapsidation sequence of human adenoviruses (in particular serotypes Ad2 and Ad5) is described in the literature, as are those of the canine adenoviruses (in
15 particular CAV1 and CAV2). In the case of the Ad5 adenovirus, for example, the encapsidation sequence corresponds to the region encompassing nucleotides 194 to 358 of the genome.

More preferably, the adenoviruses of the
20 invention encompass the ITR sequences and an encapsidation sequence, and possess a deletion of all or part of the E1 and E4 genes.

In a particularly preferred embodiment, all or part of the E1, E3 and E4 genes and, even more
25 preferably, all or part of the E1, E3, L5 and E4 genes are deleted from the genome of the adenoviruses according to the invention.

The adenoviruses of the invention can be

prepared from adenoviruses of varying origin. Thus, different serotypes of adenovirus exist whose structures and properties vary to some extent but which exhibit a comparable genetic organization. Consequently, the teaching described in the present application can easily be reproduced by the person skilled in the art for any type of adenovirus.

More specifically, the adenoviruses of the invention can be of human, animal or mixed (human and animal) origin.

As regards adenoviruses of human origin, preference is given to using those which are classed within the C group. More preferably, preference is given, among the different serotypes of human adenovirus, to using, within the scope of the present invention, type 2 or type 5 (Ad 2 or Ad 5) adenoviruses.

As indicated above, the adenoviruses of the invention can also be of animal origin or include sequences which are derived from adenoviruses of animal origin. Thus, the Applicant has demonstrated that adenoviruses of animal origin are able to infect human cells in a highly efficient manner and that they are unable to propagate themselves in the human cells in which they have been tested (cf. Application FR 93 05954). The Applicant has also demonstrated that the adenoviruses of animal origin are in no way trans-complemented by adenoviruses of human origin, thereby eliminating any risk of recombination and propagation in

vivo in the presence of a human adenovirus, which may lead to formation of an infectious particle. The use of adenoviruses or of adenovirus regions of animal origin is therefore particularly advantageous since the risks which
5 are inherent in the use of viruses as vectors in gene therapy are even lower.

The adenoviruses of animal origin which can be used within the scope of the present invention can be of canine, bovine, murine (example: Mav 1, Beard et al.,
10 Virology 75 (1990) 81), ovine, porcine, avian or else simian (example: SAV) origin. More specifically, those avian adenoviruses which may be mentioned are serotypes 1 to 10 which are available from the ATCC, such as, for example, the Phelps (ATCC VR-432), Fontes (ATCC VR-280),
15 P7-A (ATCC VR-827), IBH-2A (ATCC VR-828), J2-A (ATCC VR-829), T8-A (ATCC VR-830) or K-11 (ATCC VR-921) strains or else the strains referenced ATCC VR-831 to 835. Those bovine adenoviruses which may be used are the different known serotypes, in particular those which are available
20 from the ATCC (types 1 to 8) under reference numbers ATCC VR-313, 314, 639-642, 768 and 769. Murine adenoviruses FL (ATCC VR-550) and E20308 (ATCC VR-528), ovine adenovirus type 5 (ATCC VR-1343) or type 6 (ATCC VR-1340), porcine adenovirus 5359, or simian adenoviruses such as, in
25 particular, the adenoviruses referenced at ATCC under numbers VR-591-594, 941-943, 195-203, etc., may also be mentioned.

Among the different adenoviruses of animal

origin, preference is given, within the scope of the invention, to using adenoviruses or adenovirus regions of canine origin, in particular all the strains of the CAV2 [Manhattan strain or A26/61 strain (ATCC VR-800), for example] adenoviruses. The canine adenoviruses have been the subject of numerous structural studies. Thus, complete restriction maps of adenoviruses CAV1 and CAV2 have been described in the prior art (Spibey et al., J. Gen. Virol 70 (1989) 165), and the E1a and E3 genes as well as the ITR sequences have been cloned and sequenced (see, in particular, Spibey et al., Virus Res. 14 (1989) 241; Linné, Virus Res. 23 (1992) 119, WO 91/11525).

The defective recombinant adenoviruses according to the invention can be prepared in different ways.

A first method consists in transfecting the DNA of the defective recombinant virus, which has been prepared in vitro (either by ligation or in plasmid form), into a competent cell line, that is a cell line which carries, in trans, all the functions which are required for complementing a defective virus. These functions are preferably integrated into the genome of the cell, thereby enabling the risks of recombination to be avoided and conferring increased stability on the cell line.

A second approach consists in co-transfecting the DNA of the defective recombinant virus, which has been prepared in vitro (either by ligation or in plasmid

form), and the DNA of a helper virus into an appropriate cell line. When this method is used, it is not necessary to have available a competent cell line which is able to complement all the defective functions of the recombinant adenovirus. This is because some of these functions are complemented by the helper virus. This helper virus should itself be defective, and the cell line then carries in trans the functions which are required for complementing it. Of the cell lines which can be used, in particular, within the scope of this second approach, those which may be mentioned, in particular, are the human embryonic kidney line 293, KB cells, Hela, MDCK and GHK cells, etc. (cf. examples).

Subsequently, the vectors which have multiplied are recovered, purified and amplified using standard molecular biological techniques.

According to one embodiment, it is possible to prepare the DNA of the defective recombinant virus carrying the appropriate deletions and the two recombinant DNAs in vitro, either by ligation or in plasmid form. As indicated above, the vectors of the invention advantageously possess a deletion of all or part of certain viral genes, in particular the E1, E3, E4 and/or L5 genes. This deletion can correspond to any type of suppression which affects the gene under consideration. It can, in particular, be a question of deletion of all or part of the coding region of said gene and/or all or part of the promoter region for trans-

cribing said gene. The deletion is generally carried out on the DNA of the defective recombinant virus by, for example, digesting with appropriate restriction enzymes and then ligating, using molecular biological techniques as illustrated in the examples. The recombinant DNAs can then be inserted into this DNA, by enzymic cleavage followed by ligation, within selected regions and in the chosen orientation.

The DNA which is thus obtained, and which consequently carries the appropriate deletions and the two recombinant DNAs, enables the defective recombinant adenovirus, carrying the said deletions and recombinant DNAs, to be generated directly. This first variant is particularly well suited for achieving recombinant adenoviruses in which the genes are arranged in the form of a single transcriptional unit, or under the control of separate promoters but inserted into the same site in the genome.

It is also possible to prepare the recombinant virus in two steps, enabling the two recombinant DNAs to be introduced successively. In this case, the DNA of a first recombinant virus, carrying the appropriate deletions (or some of said deletions), and one of the recombinant DNAs is constructed, by ligation or in plasmid form. This DNA is then used to generate a first recombinant virus which carries said deletions and one recombinant DNA. The DNA of this first virus is then isolated and co-transfected with a second plasmid or the

DNA of a second defective recombinant virus which carries the second recombinant DNA, the appropriate deletions (that part not present on the first virus) and a region permitting homologous recombination. This second step
5 thereby generates the defective recombinant virus carrying the two recombinant DNAs. This preparation variant is particularly suitable for preparing recombinant viruses which carry two recombinant DNAs which are inserted into two different regions of the
10 genome of the adenovirus.

The two agents according to the invention, namely the immunosuppressant and the recombinant adenovirus, can be formulated with a view to administering them by any of the topical, oral,
15 parenteral, intranasal, intravenous, intramuscular, subcutaneous, intraocular, transdermal, etc. routes.

Preferably, the respective pharmaceutical formulation(s) contain(s) excipients which are pharmaceutically acceptable for an injectable
20 formulation. These excipients can, in particular, be sterile, isotonic salt solutions (monosodium or disodium phosphate, sodium, potassium, calcium or magnesium chloride, etc., or mixtures of such salts), or dry, in particular lyophilized, compositions which, by adding, as
25 the case may be, sterilized water or physiological saline, enable injectable solutions to be constituted.

The doses of immunosuppressant and of adenovirus which are used for the injection can be

adapted in accordance with different parameters, in particular in accordance with the mode of administration which is used, the pathology concerned, the gene to be expressed, or else the sought-after duration of the treatment.

In a general manner, the recombinant adenoviruses according to the invention are formulated and administered in the form of doses containing between 10^4 and 10^{14} pfu/ml, preferably from 10^6 to 10^{10} pfu/ml. The term pfu ("plaque-forming unit") corresponds to the infective power of a solution under consideration and is determined by infecting a suitable cell culture and measuring, generally after 5 days, the number of plaques of infected cells. The techniques for determining the pfu titre of a viral solution are well documented in the literature. As far as the immunosuppressants, more specifically, are concerned, their doses and modes of injection vary in accordance with their nature. Adjustment of these two parameters comes within the competence of the person skilled in the art.

The medicinal combination according to the invention can be used for treating or preventing numerous pathologies. Depending on the therapeutic gene which is inserted into its adenovirus, it can be used, in particular, for treating or preventing genetic disorders (dystrophy, mucoviscidosis, etc.), neurodegenerative diseases (Alzheimer's, Parkinson's, ALS, etc.), hyperproliferative pathologies (cancers, restenosis,

etc.), pathologies associated with coagulation disorders or with dyslipoproteinaemias, pathologies associated with viral infections (hepatitis, AIDS, etc.), etc.

The present invention also relates to any
 5 method of therapeutic treatment which employs the claimed medicinal combination.

The present invention will be more completely described with the aid of the examples which follow and which should be considered as being illustrative and not
 10 limiting.

Figure 1: Genetic organization of the Ad5 adenovirus. The complete sequence of Ad5 is available on database and enables the person skilled in the art to select or create any restriction site and thus to isolate any region of
 15 the genome.

Figure 2: Restriction map of the Manhattan strain of the CAV2 adenovirus (according to Spibey et al. cited above).

Figure 3: Construction of the vector pAD5-gp19k- β gal.

Figure 4: Construction of the adenovirus
 20 Ad-gp19k- β gal, Δ E1, Δ E3.

General molecular biological techniques

The methods which are routinely used in molecular biology, such as preparative extractions of plasmid DNA, centrifugation of plasmid DNA in a caesium
 25 chloride gradient, electrophoresis on agarose or acrylamide gels, purification of DNA fragments by electroelution, extraction of proteins with phenol or

with phenol/chloroform, precipitation of DNA in a saline medium with ethanol or with isopropanol, transformation into *Escherichia coli*, etc. are well known to the person skilled in the art and are amply described in the literature [Maniatis T. et al., "Molecular Cloning, A Laboratory Manual", Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1982; Ausubel F.M. et al. (eds), "Current Protocols in Molecular Biology", John Wiley & Sons, New York, 1987].

10 The plasmids of the pBR322 and pUC type, and the phages of the M13 series, were obtained commercially (Bethesda Research Laboratories).

15 For the ligations, the DNA fragments can be separated according to their size by electrophoresis in agarose or acrylamide gels, extracted with phenol or with a phenol/chloroform mixture, precipitated with ethanol and then incubated in the presence of phage T4 DNA ligase (Biolabs) in accordance with the supplier's recommendations.

20 The protruding 5' ends can be filled in using the Klenow fragment of *E. coli* DNA polymerase I (Biolabs) in accordance with the supplier's specifications. The protruding 3' ends are destroyed in the presence of phage T4 DNA polymerase (Biolabs) which is used in accordance with the manufacturer's recommendations. The protruding 25 5' ends are destroyed by carefully treating with S1 nuclease.

 In vitro site-directed mutagenesis using

synthetic oligodeoxynucleotides can be carried out using the method developed by Taylor et al. [Nucleic Acids Res. 13 (1985) 8749-8764] and employing the kit distributed by Amersham.

5 Enzymic amplification of DNA fragments by means of the technique termed PCR [polymerase-catalyzed chain reaction, Saiki R.K. et al., Science 230 (1985) 1350-1354; Mullis K.B. and Faloona F.A., Meth. Enzym. 155 (1987) 335-350] can be carried out using a DNA thermal
10 cycler (Perkin Elmer Cetus) in accordance with the manufacturer's specifications.

 The nucleotide sequences can be verified by means of the method developed by Sanger et al. [Proc. Natl. Acad. Sci. USA, 74 (1977) 5463-5467] using the kit
15 distributed by Amersham.

Cell lines employed

 In the examples which follow, the following cell lines have been or can be employed:

- Human embryonic kidney line 293 (Graham et
20 al., J. Gen. Virol. 36 (1977) 59). This line contains in particular, integrated into its genome, the left-hand part of the genome of the human adenovirus Ad5 (12%).
- KB human cell line. Derived from a human epidermal carcinoma, this line can be obtained from ATCC
25 (ref. CCL17) as can the conditions for culturing it.

- HeLa human cell line: derived from a human epithelium carcinoma, this line can be obtained from ATCC (ref. CCL2) as can the conditions for culturing it.

- MDCK canine cell line: the conditions for
5 culturing MDCK cells have been described, in particular, by Macatney et al., Science 44 (1988) 9.

- gm DBP6 cell line (Brough et al., Virology 190 (1992) 624). This line consists of HeLa cells carrying the adenovirus E2 gene under the control of the
10 LTR of MMTV.

EXAMPLES

Example 1. Construction of defective recombinant adenoviruses encompassing a therapeutic gene (the LacZ gene of E. coli) under the control of the LTR promoter of RSV and
15 the gp19k gene under the control of the LTR promoter of RSV, with both genes being inserted within the E1 region.

These adenoviruses were constructed by homologous recombination between a plasmid carrying the left-hand part of the Ad5 adenovirus, the two recombinant
20 DNAs and a region of the Ad5 adenovirus (corresponding to protein IX) and the DNA of a defective adenovirus carrying various deletions.

1. Construction of the vector pAD5-gp19k- β gal (Figure 3)

25 1.1. Construction of the plasmid pGEM-gp19k

Plasmid pAD5-gp19k- β gal contains a cDNA sequence encoding the adenovirus protein gp19k. This plasmid was constructed as follows. The XbaI fragment of the genome of wild-type Ad5 adenovirus, containing the E3 region, was isolated and cloned into the corresponding site of plasmid pGEM (Promega) in order to generate plasmid pGEM-E3. The HinfI fragment, containing the sequence encoding gp19k (nucleotides 28628 to 29634 of wild-type Ad5 adenovirus), was then isolated from plasmid pGEM-E3. The ends of this fragment were rendered blunt by the action of the Klenow fragment of *E. coli* DNA polymerase I (cf. general molecular biological techniques) and the fragment which was obtained was then cloned into the SmaI site of plasmid pGEMzf+ (Promega).

The plasmid which was obtained was designated pGEM-gp19k (Figure 3).

1.2. Construction of the vector pAD5-gp19k- β gal

This example describes the construction of a plasmid which contains one of the two recombinant DNAs encompassing their own promoter, the left-hand part of the adenovirus genome and a supplementary part (protein pIX) permitting homologous recombination. This vector was constructed from the plasmid pAd.RSV β Gal as follows.

The plasmid pAd.RSV β Gal contains, in the 5'>3' orientation,

- the PvuII fragment corresponding to the left-hand end of adenovirus Ad5 encompassing: the ITR

sequence, the origin of replication, the encapsidation signals and the E1A enhancer,

- the gene encoding β -galactosidase under the control of the RSV promoter (from Rous sarcoma virus).

5 - a second fragment of the genome of adenovirus Ad5, which permits homologous recombination between plasmid pAd.RSV β Gal and the adenovirus d1324. Plasmid pAd.RSV β Gal has been described by Stratford-Perricaudet et al. (J. Clin. Invest. 90 (1992) 626).

10 Plasmid pAd.RSV β Gal was first of all cut with the enzymes EagI and ClaI. This generates a first fragment carrying, in particular, the left-hand part of adenovirus Ad5 and the LTR promoter from RSV. In parallel, the plasmid pAd.RSV β Gal was also cut with the
15 enzymes EagI and XbaI. This generates a second type of fragment carrying, in particular, the LTR promoter of RSV, the LacZ gene and a fragment of the genome of adenovirus Ad5 which permits homologous recombination. The ClaI-EagI and EagI-XbaI fragments were then ligated
20 in the presence of the XbaI-ClaI fragment from plasmid pGEM-gp19k (Example 1.1) carrying the sequence encoding gp19k (cf. Figure 3). The vector which was obtained in this way, designated pAD5-gp19k- β gal, therefore contains

- the PvuII fragment corresponding to the left-
25 hand end of adenovirus Ad5 encompassing: the ITR sequence, the origin of replication, the encapsidation signals and the E1A enhancer,

- the sequence encoding gp19k under the control

of the RSV promoter (from Rous sarcoma virus),

- the gene encoding β -galactosidase under the control of the RSV promoter (from Rous sarcoma virus), and

5 - a second fragment of the genome of adenovirus Ad5 which permits homologous recombination.

2. Construction of the recombinant adenoviruses

2.1. Construction of a recombinant adenovirus which is deleted in the E1 region and which carries the
10 two recombinant DNAs inserted in the same orientation within the E1 region.

Vector pAD5-gp19k- β gal was linearized and cotransfected with an adenoviral vector, which was deficient in the E1 gene, into helper cells (line 293)
15 which supplied in trans the functions encoded by the adenovirus E1 (E1A and E1B) regions.

More precisely, the adenovirus Ad-gp19k- β gal, Δ E1 is obtained by homologous recombination in vivo between the adenovirus Ad-RSV β gal (cf. Stratford-
20 Perricaudet et al. cited above) and vector pAD5-gp19k- β gal in accordance with the following protocol: plasmid pAD5-gp19k- β gal, which is linearized with XmnI, and adenovirus Ad-RSV β gal, which is linearized with the enzyme ClaI, are co-transfected into line 293 in the
25 presence of calcium phosphate in order to enable homologous recombination to take place. The recombinant adenoviruses which are generated in this way are then

selected by plaque purification. Following isolation, the DNA of the recombinant adenovirus is amplified in cell line 293, resulting in a culture supernatant which contains the unpurified defective recombinant adenovirus with a titre of approximately 10^{10} pfu/ml.

In general, the viral particles are purified by centrifugation in a caesium chloride gradient in accordance with known techniques (see, in particular, Graham et al., Virology 52 (1973) 456). The adenovirus Ad-gp19k- β gal, Δ E1 can be stored at -80°C in 20% glycerol.

2.2 Construction of a recombinant adenovirus which is deleted in the E1 and E3 regions and which carries the two recombinant DNAs inserted in the same orientation within the E1 region (Figure 4).

Vector pAD5-gp19k- β gal was linearized and cotransfected with an adenoviral vector, which was deficient in the E1 and E3 genes, into helper cells (line 293) which supply in trans the functions encoded by the adenovirus E1 (E1A and E1B) regions.

More precisely, the adenovirus Ad-gp19k- β gal, Δ E1, Δ E3 was obtained by homologous recombination in vivo between the mutant adenovirus Ad-dl1324 (Thimmappaya et al, Cell 31 (1982) 543) and vector pAD5-gp19k- β gal in accordance with the following protocol: plasmid pAD5-gp19k- β gal and adenovirus Ad-dl1324, linearized with the enzyme ClaI, were cotransfected into line 293 in the presence of calcium phosphate in order to enable homologous recombination to take place. The recombinant

adenoviruses which were generated in this way were then selected by plaque purification. Following isolation, the DNA of the recombinant adenovirus is amplified in cell line 293, resulting in a culture supernatant which contains the unpurified defective recombinant adenovirus with a titre of approximately 10^{10} pfu/ml.

In general, the viral particles are purified by centrifugation in a caesium chloride gradient in accordance with known techniques (see, in particular, Graham et al. Virology 52 (1973) 456). The genome of the recombinant adenovirus was then verified by Southern blot analysis. Adenovirus Ad-gp19k- β gal, Δ E1, Δ E3 can be stored at -80°C in 20% glycerol.

Example 2: Demonstration of the immunoprotective activity of the medicinal combination according to the invention.

60 adult female DBA/2 mice are divided randomly into 6 groups of 10 mice which are then treated respectively in accordance with the following injection protocols:

20 - GROUP 1a:

Is given an intraocular injection of 10 μg of anti-CD3 monoclonal antibodies on days -2, -1, 1, 2, 3, 4 and 5 with an intravenous injection of $4 \cdot 10^9$ pfu of Ad-RSV β gal virus on day 0 (cf. Stratford-Perricaudet et al. cited above).

- GROUP 1b:

Is given the same treatment as group 1a but

employing, as virus, 4.10^9 pfu of Ad-gp 19k- β gal virus (Figure 4).

- GROUP 2a:

Is given an intraperitoneal injection of 250 μ g
5 of anti-CD4 monoclonal antibodies on days -2, -1, 1, 4
and 7 with an intravenous injection of 4.10^9 pfu of Ad-RSV
 β gal virus on day 0.

- GROUP 2b:

Is given the same treatment as group 2a but
10 using, as virus, 4.10^9 pfu of Ad gp 19k- β g al virus.

- GROUP 3a:

Is given an intravenous injection of 4.10^9 pfu
of Ad- β gal without any accompanying administration of
immunosuppressant.

15 - GROUP 3b:

Is given an intravenous injection of 4.10^9 pfu
of Ad-gp19k- β gal without any accompanying administration
of immunosuppressant.

At various times, two animals from each group
20 were sacrificed with the aim of removing their livers and
spleens.

2.1 - Immunofluorescence analysis of the
distribution of the principal lymphocyte subpopulations
(CD3+, CD4+ and CD8+) within splenocytes which are
25 removed on D15 after the injection.

A suspension of isolated cells was prepared
from removed spleens. A cell sample was analysed by
immunofluorescence using antibodies which were specific

for each lymphocyte subpopulation. The fluorescent cells were read with the aid of a cytofluorimeter (Becton Dickinson FACS Scan). The results are given in Table I below.

	Group 3a Ad- β gal	Group 3b Ad- β gal- gp19k		Group 1a anti-CD3/ Ad- β gal	Group 2a anti-CD4/ Ad/ β gal				
	% of cells expressing β gal at the cell surface								
5	CD3	20.0	17.5	20.6	21	5.4	6.1	12	10.3
	CD4	13.4	12.6	15.3	16.8	4.4	5.1	2.7	4.1
	CD8	5.5	5.5	6.1	6	2.02	2.3	7.9	6.6

TABLE I

The clear decrease in the CD3+, CD4+ and CD8+ cells in the animals treated with anti-CD3 is noted as is the selective decrease in CD4+ cells in the animals treated with anti-CD4.

2.2. - Analysis of the cytotoxic capacity of the splenocytes which are removed at D32 after the injection and stimulated in vitro with regard to histocompatible target cells expressing β gal

A second splenocyte sample isolated from the spleens of treated animals was cultured in vitro for 4

days in the presence of P815 cells expressing β -galactosidase at their surface. At the end of the culture, the cytotoxic activity of these splenocytes with regard to P815- β gal target cells labelled with Cr^{51} was evaluated. The cytotoxic activity, expressed as per cent cytolysis, was determined in a conventional manner by bringing together different ratios of effector cells and target cells. The results are presented in Table II below.

10	Group	2b	3A
	Treatment	Anti-CD4/ Ad- β gal-gp19k	AD- β gal
	Ratio Effector/targets	% cytolysis	
	80/1	4 2	13 14
15	40/1	2 1	13 9
	20/1	1 1	5 9
	10/1	0 0	2 5
	5/1	0 1	1 2

There is seen to be a very clear neutralization of the cytotoxic capacity of the splenocytes which were removed from the animals having been treated with anti-

CD4, that is to say the group 2b.

TABLE II

2.3. Expression of β -galactosidase activity in the liver after 15 and 32 days.

5 The livers are sectioned and stained with X-gal in order to display the β -galactosidase activity and with eosin in order to demonstrate the histology of the section. The results are presented in Table III below.

	<u>Number of cells expressing βgal</u>	
	15 days	32 days
10 Group 2a: (anti-CD4/Ad- β gal)	1	1
Group 2b: (anti-CD4/Ad gp19k- β gal)	250	50
Group 3a: (Ad- β gal)	3	0
15 Group 3b: (Ad gp19k- β gal)	25	0

TABLE III

From the results presented above, it emerges that injection of anti-CD4 antibodies in association with

an injection of Adgp19k- β gal induces an expression of the gene under consideration which is markedly prolonged. Thus, 30 days after the injections, significant β -galactosidase activity is observed in the case of group 5 2b. This prolongation, which can be interpreted as the result of a tolerance phenomenon which is induced in accordance with the invention, is markedly greater than that which could have been expected from the simple juxtaposition of the respective effects of the anti-CD4 10 immunosuppressants and of the recombinant adenovirus Ad gp19k- β gal.

Furthermore, no inflammatory reaction is observed over this period of 30 days in the case of group 2b.

CLAIMS

1. Medicinal combination of at least one immunosuppressive agent and at least one recombinant adenovirus whose genome comprises a first recombinant DNA containing a therapeutic gene and a second recombinant DNA containing an immunoprotective gene, for consecutive, intermittent and/or simultaneous use over time, which can be used for exogenous transfections in vivo and/or ex vivo.
2. Medicinal combination according to claim 1, characterized in that the immunosuppressive agent is preferably selected from among cyclosporin, FK506, azathioprine, corticosteroids and monoclonal or polyclonal antibodies.
3. Medicinal combination according to claim 2, characterized in that the antibodies concerned are antibodies which are able to inactivate immune molecules or induce destruction of the immune cells carrying these molecules.
4. Medicinal combination according to claim 3, characterized in that the antibody is selected from among the anti-CD4, -CD2, -CD3, -CD8, -CD28, -B7, -ICAM-1 and -LFA-1 antibodies and CTLA4Ig.
5. Medicinal combination according to one of the preceding claims, characterized in that the therapeutic gene encodes a therapeutic protein.
6. Medicinal combination according to one of claims 1 to 4, characterized in that the therapeutic gene

inserted in opposite orientations.

14. Medicinal combination according to one of the preceding claims, characterized in that the two recombinant DNAs are inserted into one and the same site
5 of the adenovirus genome, preferably within the E1, E3 or E4 regions.

15. Medicinal combination according to claim 14, characterized in that the two recombinant DNAs are inserted within the E1 region.

10 16. Medicinal combination according to one of claims 1 to 13, characterized in that the two recombinant DNAs are inserted into different sites in the adenovirus genome.

15 17. Medicinal combination according to claim 16, characterized in that one of the recombinant DNAs is inserted within the E1 region and the other within the E3 or E4 region.

20 18. Medicinal combination according to one of the preceding claims, characterized in that the adenovirus is a defective recombinant adenovirus which encompasses the ITR sequences and a sequence permitting encapsidation and which carries a deletion of all or part of the E1 and E4 genes.

25 19. Medicinal combination according to claim 18, characterized in that the adenovirus concerned is an adenovirus which encompasses the ITR sequences and a sequence permitting encapsidation and which carries a deletion of all or part of the E1, E3 and E4 genes.

20. Medicinal combination according to one of claims 1 to 19, characterized in that the adenovirus concerned is an adenovirus from whose genome all or part of the E1, E3, L5 and E4 genes have been deleted.

5 21. Medicinal combination according to one of the preceding claims, characterized in that the recombinant adenovirus is of human, animal or mixed origin.

10 22. Medicinal combination according to claim 21, characterized in that the recombinant adenoviruses of human origin are selected from among those classed within the C group, preferably from among the type 2 or type 5 recombinant adenoviruses (Ad 2 or Ad 5).

15 23. Medicinal combination according to claim 22, characterized in that the adenoviruses of animal origin are chosen from among the adenoviruses of canine, bovine, murine, ovine, porcine, avian and simian origin.

20 24. Medicinal combination according to one of the preceding claims, characterized in that the immunosuppressive agent is injected before and after injection of the adenovirus.

25 25. Medicinal combination according to one of the preceding claims, characterized in that the immunosuppressive agent and the recombinant adenovirus are injected simultaneously.

5

Abstract

A medical combination of at least one immunosuppressive
10 agent and at least one recombinant adenovirus with a genome that
includes a first recombinant DNA containing a therapeutic gene,
and a second recombinant DNA containing an immunoprotective
gene, for consecutive, intermittent and/or simultaneous use in
in vivo and/or *ex vivo* exogenic transfections.

"Abstract" sheet







Figure 1





4/4



**DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION**

As a below named inventor, and on behalf of co-inventor Martin Lee (64 Princes Street, Oxford, England, Citizenship British), I hereby declare that my residence, post office address and citizenship are as stated below next to my name; I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

**MEDICINAL COMBINATION USEFUL FOR IN VIVO EXOGENIC
TRANSFECTION AND EXPRESSION**

the international specification of which was filed on **February 12, 1996** as Application Serial No. **PCT/FR96/00218** which notice of transmission was given on **August 22, 1996**, by the International Bureau. I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a). I hereby claim foreign priority benefits under Title 35, United States Code, Section 119 of a foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Applications(s)			Priority Claimed	
<u>FR95/01662</u>	<u>France</u>	<u>14 February 1995</u>	<u>X</u>	<u> </u>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No
<u> </u>	<u> </u>	<u> </u>	<u>Yes</u>	<u>No</u>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56(a) which occurred between the filing date of the prior application and the national or PCT International filing date of this application:

<u>(Application Serial No.)</u>	<u>(Filing Date)</u>	<u>(Status-Patented, Pending or Abandoned)</u>
---------------------------------	----------------------	--

<u>(Application Serial No.)</u>	<u>(Filing Date)</u>	<u>(Status-Patented, Pending or Abandoned)</u>
---------------------------------	----------------------	--

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: I (We) hereby appoint the attorneys associated with the Customer Number provided below as my (our) attorneys, with full powers of substitution and revocation, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

Customer No.: 005487
Direct Telephone Calls to: (610) 454-3839

Michel PERRICAUDET	French
First or Sole Inventor	Citizenship

31 rue de Chartres	Same
Residence Address	Post Office Address

28320 Ecrosnes	FLX	Same
City		City

FRANCE	Same
State (Zip) or Country	State (Zip) or Country

Date May 5th, 1998 Signature Michel Perricaudet

<u>Lucienne CHATENOUD</u>	French
Second Inventor	Citizenship

160 Avenue de Versailles	Same
Residence Address	Post Office Address

75016 Paris	Same
City	City

FRANCE	Same
State (Zip) or Country	State (Zip) or Country

Date _____ Signature _____

2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2809 2810 2811 2812 2813 2814 2815 2816 2817 2

200

Direct Telephone Calls to: (610) 454-3839

Date
Signature

Date _____ Signature _____

463 Hedi HADDADA

Third Inventor

French

Citizenship

18 Avenue de Bellevue

Residence Address

Same

Post Office Address

92340 Bg la Reive

City

FRX

Same

City

FRANCE

State (Zip) or Country

Same

State (Zip) or Country

Date

05/5/98

Signature

Haddada

Jean-François BACH

Fourth Inventor

French

Citizenship

180 rue de Grenelle

Residence Address

Same

Post Office Address

75007 Paris

City

Same

City

FRANCE

State (Zip) or Country

Same

State (Zip) or Country

Date

Signature

Michelle WEBB

Fifth Inventor

British

Citizenship

125 Broomwood Road

Residence Address

Same

Post Office Address

London SW11 6JU

City

Same

City

ENGLAND

State (Zip) or Country

Same

State (Zip) or Country

Date

Signature

**DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION**

As a below named inventor, and on behalf of co-inventor Martin Lee (64 Princes Street, Oxford, England, Citizenship British), I hereby declare that my residence, post office address and citizenship are as stated below next to my name; I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

**MEDICINAL COMBINATION USEFUL FOR IN VIVO EXOGENIC
TRANSFECTION AND EXPRESSION**

the international specification of which was filed on February 12, 1996 as Application Serial No. PCT/FR96/00218 which notice of transmission was given on August 22, 1996, by the International Bureau. I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a). I hereby claim foreign priority benefits under Title 35, United States Code, Section 119 of a foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Applications(s)			Priority Claimed	
<u>FR95/01662</u>	<u>France</u>	<u>14 February 1995</u>	<u>X</u>	<u> </u>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No
<hr/>				
<u> </u>	<u> </u>	<u> </u>	<u> </u>	<u> </u>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56(a) which occurred between the filing date of the prior application and the national or PCT International filing date of this application:

<u> </u>	<u> </u>	<u> </u>
(Application Serial No.)	(Filing Date)	(Status-Patented, Pending or Abandoned)
<hr/>		
<u> </u>	<u> </u>	<u> </u>
(Application Serial No)	(Filing Date)	(Status-Patented, Pending or Abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: I (We) hereby appoint the attorneys associated with the Customer Number provided below as my (our) attorneys, with full powers of substitution and revocation, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

Customer No.: 005487
Direct Telephone Calls to: (610) 454-3839

<u>Michel PERRICAUDET</u>	<u>French</u>
First or Sole Inventor	Citizenship
<u>31 rue de Chartres</u>	<u>Same</u>
Residence Address	Post Office Address
<u>28320 Ecrosnes</u>	<u>Same</u>
City	City
<u>FRANCE</u>	<u>Same</u>
State (Zip) or Country	State (Zip) or Country
<u> </u>	<u> </u>
Date	Signature

=====

<u>300 Lucienne CHATENOU</u>	<u>French</u>
Second Inventor	Citizenship
<u>160 Avenue de Versailles</u>	<u>Same</u>
Residence Address	Post Office Address
<u>75016 Paris</u> <u>FR X</u>	<u>Same</u>
City	City
<u>FRANCE</u>	<u>Same</u>
State (Zip) or Country	State (Zip) or Country
<u>Date 6/5/98</u>	<u>Signature</u>

=====

Date _____ Signature _____

Date _____ Signature _____

Date 12.4.98 Signature MC Webb

1224248

Atkinson

[illegible]

**DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION**

As a below named inventor, and on behalf of co-inventor Martin Lee (64 Princes Street, Oxford, England, Citizenship British), I hereby declare that my residence, post office address and citizenship are as stated below next to my name; I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

**MEDICINAL COMBINATION USEFUL FOR IN VIVO EXOGENIC
TRANSFECTION AND EXPRESSION**

the international specification of which was filed on February 12, 1996 as Application Serial No. PCT/FR96/00218 which notice of transmission was given on August 22, 1996, by the International Bureau. I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a). I hereby claim foreign priority benefits under Title 35, United States Code, Section 119 of a foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Applications(s)			Priority Claimed	
<u>FR95/01662</u>	<u>France</u>	<u>14 February 1995</u>	<u>X</u>	<u> </u>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No
<hr/>			<hr/>	<hr/>
(Number)	(Country)	(Day/Month/Year Filed)	Yes	No

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56(a) which occurred between the filing date of the prior application and the national or PCT International filing date of this application:

<u> </u>	<u> </u>	<u> </u>
(Application Serial No.)	(Filing Date)	(Status-Patented, Pending or Abandoned)
<hr/>		
<u> </u>	<u> </u>	<u> </u>
(Application Serial No.)	(Filing Date)	(Status-Patented, Pending or Abandoned)

and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: I (We) hereby appoint the attorneys associated with the Customer Number provided below as my (our) attorneys, with full powers of substitution and revocation, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

Customer No.: 005487

Direct Telephone Calls to: (610) 454-3839

Michel PERRICAUDET	French
First or Sole Inventor	Citizenship
31 rue de Chartres	Same
Residence Address	Post Office Address
28320 Ecrosnes	Same
City	City
FRANCE	Same
State (Zip) or Country	State (Zip) or Country

Date

Signature

=====

Lucienne CHATENOUD	French
Second Inventor	Citizenship
160 Avenue de Versailles	Same
Residence Address	Post Office Address
75016 Paris	Same
City	City
FRANCE	Same
State (Zip) or Country	State (Zip) or Country

Date

Signature

=====

